

**COMPENDIUM OF  
VETERINARY STANDARD PRECAUTIONS:  
ZOO NOTIC DISEASE PREVENTION IN VETERINARY  
PERSONNEL**

**National Association of State Public Health Veterinarians  
Veterinary Infection Control Committee  
2006**

# **Compendium of Veterinary Standard Precautions: Zoonotic Disease Prevention in Veterinary Personnel, 2006**

**National Association of State Public Health Veterinarians (NASPHV)  
Veterinary Infection Control Committee (VICC)**

## **SUMMARY**

Veterinary practices are unique environments that bring humans into close contact with many different species of ill animals. In the practice environment, whether in a building or “in the field,” veterinary personnel are frequently exposed to recognized and unrecognized infectious pathogens, many of which are zoonotic (transmitted from animals). Some of the documented zoonotic infections that have occurred in veterinary personnel include: multi-drug resistant *Salmonella* Typhimurium, cryptosporidiosis, cat-associated plague, cat-associated sporotrichosis, Methicillin-resistant *Staphylococcus aureus* (MRSA), and dermatophytosis.

Infection control measures vary from practice to practice and are often insufficient to prevent zoonotic disease transmission. The Veterinary Standard Precautions (VSP) in this compendium are designed to prevent transmission of zoonotic pathogens from animal patients to veterinary personnel in private practice. The compendium committee has set the following objectives:

- Raise awareness of the scope of zoonotic disease risk in veterinary practice
- Address issues specific to veterinary practices, such as bites and other trauma from veterinary patients, and housing and monitoring of ill animals in common treatment areas
- Provide practical, science-based guidance that will reduce transmission from recognized and unrecognized sources of infection
- Outline sensible infection control practices
- Provide a model infection control plan for use by individual veterinary practices

### **The NASPHV Committee**

Brigid Elchos, RN, DVM , CoChair

Joni Scheftel, DVM, MPH, DACVPM, CoChair

Bryan Cherry, VMD, PhD

Emilio DeBess, DVM, MPVM

Sharon Hopkins, DVM, MPH

Lee Hunter, DVM, MPH

Jay Levine, DVM, MPH

Carl Williams, DVM, MA

**Consultants to the Committee**

Mike Bell, MD: Association for Professionals in Infection Control and Epidemiology (APIC) and Centers for Disease Control and Prevention (CDC)

Boris Pavlin, MD: CDC

Glenda Dvorak, DVM, MPH: Center for Food Security and Public Health, IA

Kathy Orloski, DVM, MS, DACVPM: USDA APHIS VS Center for Emerging Issues

Rebecca Stinson-Dixon, DVM: American Veterinary Medical Association (AVMA)

Heidi Burnett, DVM: American Animal Hospital Association (AAHA)

Jo Hofmann, MD: Council of State and Territorial Epidemiologists (CSTE)

Ryan Cheek, RVT, VTS (ECC): National Association of Veterinary Technicians in America (NAVTA)

**Address correspondence to:** Dr. Joni Scheftel, Co-chair, NASPHV VSP Compendium Committee, Acute Disease Investigation and Control, Minnesota Department of Health, 625 Robert St. N., P.O. Box 64975, St. Paul, MN 55164-0975. Telephone: 651-201-5107; Fax: 651-201-5743; E-mail: [joni.scheftel@health.state.mn.us](mailto:joni.scheftel@health.state.mn.us)

## **TABLE OF CONTENTS**

### **I. INTRODUCTION**

- A. Background and Objectives**
- B. Considerations**

### **II. ZOO NOTIC DISEASE TRANSMISSION**

- A. Transmission Mechanisms**
  - 1. Contact**
  - 2. Aerosol**
  - 3. Vector-borne**

### **III. VETERINARY STANDARD PRECAUTIONS**

- A. Personal Protective Actions and Equipment**
  - 1. Hand Hygiene**
  - 2. Use of Gloves and Sleeves**
  - 3. Facial Protection**
  - 4. Respiratory Protection**
  - 5. Protective Outerwear**
    - a. Lab Coats, Smocks, and Coveralls**
    - b. Non-Sterile Gowns**
    - c. Footwear**
    - d. Headcovers**
  - 6. Bite and Other Animal-Related Injury Prevention**
- B. Protective Actions During Veterinary Procedures**
  - 1. Intake**
  - 2. Examination of Animals**
  - 3. Injections, Venipuncture, Aspirations**
    - a. Needlestick Injury Prevention**
    - b. Barrier Protection**
  - 4. Dental Procedures**
  - 5. Resuscitation**
  - 6. Obstetrics**
  - 7. Necropsy**
  - 8. Diagnostic Specimen Handling**
- C. Environmental Infection Control**
  - 1. Isolation of Infectious Animals**
  - 2. Cleaning and Disinfection of Equipment and Environmental Surfaces**
  - 3. Handling of Laundry**
  - 4. Decontamination and Spill Response**
  - 5. Veterinary Medical Waste**
  - 6. Rodent and Vector Control**
  - 7. Other Environmental Controls**

#### **IV. EMPLOYEE HEALTH**

##### **A. General**

- 1. Infection Control Personnel**
- 2. Employee Immunization Policies and Record Keeping**
  - a. Rabies**
  - b. Tetanus**
  - c. Seasonal Influenza**
  - d. Immunization and Other Health Records**
- 3. Staff Training and Education**
- 4. Documenting Exposure Incidents**

##### **B. Immunocompromised Personnel**

#### **V. CREATING A WRITTEN INFECTION CONTROL PLAN**

##### **A. Communicating and Updating the Infection Control Plan**

- 1. Availability**
- 2. Leadership**
- 3. New Staff**
- 4. Continuing Education**
- 5. Review and Revision**
- 6. Assurance**

#### **VI. REFERENCES**

**APPENDIX A: Zoonotic Diseases of Importance in the United States**

**APPENDIX B: Selected Disinfectants Used in Veterinary Practice**

**APPENDIX C: OSHA Respiratory Protection Standard**

**APPENDIX D: Model Infection Control Plan for Veterinary Practices**

# **Compendium of Veterinary Standard Precautions: Zoonotic Disease Prevention in Veterinary Personnel 2006**

## **National Association of State Public Health Veterinarians Veterinary Infection Control Committee**

### **I. INTRODUCTION**

#### **A. Background and Objectives**

Veterinary personnel are at risk of contracting infections from animals (zoonotic diseases) because of their frequent contact with a wide variety of species (1). Yet, the scope of zoonotic disease risk has not been well recognized, and uniform guidance for infection control in general veterinary practice has been limited. Infection control practices vary tremendously from one veterinary practice to another and often are not sufficient to prevent zoonotic disease transmission (2, 3, J. McQuiston personal communication, Aug 22, 2006, J. Wright personal communication, Aug 22, 2006).

The Veterinary Standard Precautions (VSP) outlined in this compendium are designed to reduce the risk of zoonotic infections among personnel in private veterinary clinics and hospitals from both recognized and unrecognized sources of infection. They are to be used with all patients, regardless of their diagnosis, when contacting blood, all body fluids, feces, exudates, non-intact skin, or mucous membranes. In addition, VSP include practices to prevent bites and other trauma induced by veterinary patients that may result in exposure to zoonotic pathogens.

Infection control in human medicine has evolved considerably in the past 30 years with the recognition of HIV and hepatitis B and C viruses. Currently, the cornerstone of human infection control is the use of “Standard Precautions” (4), which provide a basis for the principles of VSP. It would be prudent for veterinary practices to apply standard infection control procedures now, rather than as a response to a serious or fatal zoonotic disease outbreak in veterinary personnel.

New infectious diseases are continually emerging (5). A recent report cataloging human disease agents found that of 1,415 agents causing disease in humans, 868 (61%) are zoonotic. Also, of the 175 pathogens defined as emerging infections, 75% are zoonotic (6). Moreover, the globalization of commerce and the ease of international travel heighten the likelihood of encountering zoonotic pathogens. The 2003 U.S. outbreak of human monkeypox in six states is a prime example: Eighteen (25%) of 71 human monkeypox cases were in veterinarians or veterinary staff who contracted the disease from contact with pet prairie dogs infected by imported African rodents (7).

While the introduction of novel pathogens, such as monkeypox, highlights the need for infection control, these practices would also protect veterinary personnel from common

pathogens. In fact, many endemic zoonotic infections occur in veterinary personnel; among those that have been documented in the literature are:

- Multidrug-resistant *Salmonella* Typhimurium outbreaks with zoonotic transmission to veterinary staff and students (8-10).
- Cryptosporidiosis among veterinary students in multiple states (11-15).
- Cat-associated plague among veterinary personnel in eight Western states (16).
- Cat-associated sporotrichosis in veterinary personnel (17-21).
- Methicillin-resistant *Staphylococcus aureus* (MRSA) transmission among veterinary personnel and equine, canine and feline patients (10, 22-25).
- Dermatophytosis (ringworm) among British veterinarians (26).

As a national coalition of veterinary public health professionals, NASPHV recognizes the need for specific guidance addressing zoonotic disease risk in veterinary practices. NASPHV strongly believes that infection control guidance for veterinary personnel should be provided primarily by veterinary personnel. Public health veterinarians are uniquely qualified to address this issue because they understand and regularly address animal diseases as they relate to human health. This resulted in the formation of the Veterinary Infection Control Committee (VICC) and development of the compendium.

The VICC has set the following objectives for the compendium:

- Raise awareness of the scope of zoonotic disease risk in veterinary practice
- Address issues specific to veterinary practices, such as bites and other trauma from veterinary patients, and housing and monitoring of ill animals in common treatment areas
- Provide practical, science-based guidance that will reduce transmission from recognized and unrecognized sources of infection
- Outline sensible infection control practices
- Provide a model infection control plan for use by individual veterinary practices

## **B. Considerations**

This Compendium focuses exclusively on protection of veterinary personnel in clinical practice. Although eliminating all risk from zoonotic pathogens in veterinary practice is not achievable, this report provides recommendations for minimizing disease and injury. The document may also be useful for preventing infectious agent transmission in facilities other than veterinary clinics and hospitals (such as shelters and kennels), but it is not intended to be inclusive for all animal facilities and practice types.

Since veterinary practices vary, these guidelines may be adapted to individual practice needs and circumstances, but should comply with federal, state, and local regulations. Modifications to these recommendations should adhere to the basic principles of infection control to prevent spread of occupational zoonoses by all routes of transmission.

Every effort has been made to compile existing scientific evidence for standard practices that will protect veterinary personnel from zoonotic diseases. As there have been few studies to test the efficacy of infection control in veterinary practice and gaps exist in the

knowledge of the epidemiology of some diseases, it is expected that the content of this publication will evolve.

Training is an essential part of understanding and implementing the guidelines. The Compendium will be most effective if each employee is aware of the relevance of the guidelines to his or her health and the health of others in the practice.

A client education program will help protect veterinary staff from zoonotic diseases, as well as protecting clients, their children, and other pets. Client education should include the importance of rabies vaccination, comprehensive internal and external parasite control, and bite prevention. Clients should be informed of zoonotic disease risks specific to their geographic area and the veterinarian promoted as a source of expert information on zoonotic disease control issues. Written educational information should be made available in clinic waiting areas and on websites.

Employers should promote safe work habits. The cost of implementing these guidelines should be compared to the potential consequences of inadequate infection control, including loss of business due to practice closure, sick leave or hospitalization of personnel, loss of credibility, and litigation (27-29).

## **II. ZOONOTIC DISEASE TRANSMISSION**

Transmission of infections requires three elements: a source of pathogens, a susceptible host, and a means of transmission for the microorganism (4). Sources include animals or environments contaminated by animals. Pathogens may be transmitted to humans directly from the animal via blood or other body fluids during diagnostic or treatment procedures. Animals may be clinically ill, asymptomatic carriers of an infectious agent, or harbor endogenous flora that are pathogenic to humans. Pathogens may also be transmitted indirectly from objects in the environment including walls, floors, counters, equipment, supplies, animal feed, and water.

Host resistance to pathogenic microorganisms varies greatly. Some persons may be immune to infection or may be able to resist colonization by an infectious agent; others exposed to the same agent may establish a commensal relationship with the infecting microorganism and become asymptomatic carriers; still others may develop clinical disease. Host factors such as age, underlying diseases, irradiation, pregnancy, and breaks in the body's first-line of defense mechanisms (intact skin, cough reflex, stomach acid) may render a host more susceptible to infection. Conversely, vaccination may reduce susceptibility to infection.

### **A. Transmission Mechanisms**

Transmission occurs through three main mechanisms: contact, aerosol, and vector-borne. The same agent may be transmitted by more than one route. Transmission is largely influenced by the stability of the pathogen, the pathogenesis of the disease it causes, and the routes by which it leaves the infected host. Different agents vary in their degree of infectivity through the various routes.



**1. Contact transmission** can occur when pathogens from animals or their environments enter the human host through three routes: ingestion, mucous membrane exposure, and cutaneous/percutaneous. Direct contact transmission may occur during activities such as examining, medicating, bathing, and handling animals. Indirect contact transmission involves contact with a contaminated intermediate object, such as occurs during cleaning cages and equipment and handling soiled laundry. Injuries from contaminated sharps, such as scalpel blades, needles, and necropsy knives, may result in exposure to live vaccines and pathogens. In addition, injury from sharps increases risk of exposure to other pathogens through direct and indirect contact (4).

**2. Aerosol transmission** can occur when pathogens from animals or their environments travel via the air and enter the human host through inhalation and/or mucous membranes. In general, risk to veterinary personnel increases with proximity to the source and the length of time over which exposure occurs. Transmission over short distances occurs when droplets created by coughing, sneezing, vocalizing, or procedures such as suctioning and bronchoscopy are propelled through the air and deposited on the host's conjunctivae, nasal or oral mucosa. Certain pathogens may remain infective over longer distances (4, 30). However, defining the infective distance is difficult because it depends on particle size, the nature of the pathogen, and environmental factors (30). Although data are not available to define specific infection risk from aerosol transmission for most pathogens, some pathogens known to be transmitted over longer distances include *Coxiella burnetii* (31-33) and *Mycobacterium bovis* (34).

**3. Vector-borne transmission** occurs when vectors such as mosquitoes, fleas, ticks, rats, and other animals transmit microorganisms. Animals may bring flea and tick vectors into contact with veterinary personnel. Veterinary personnel working in outdoor settings may be at risk for diseases carried by arthropods and other biological vectors.

### **III. VETERINARY STANDARD PRECAUTIONS**

#### **A. Personal Protective Actions and Equipment**

##### **1. Hand Hygiene**

Handwashing is the single most important measure to reduce the risk of disease transmission (4, 35, 36). Hands should be washed between animal contacts and after contact with blood, body fluids, secretions, excretions, and equipment or articles contaminated by them. Staff with animal contact should keep fingernails short (37).

Correct handwashing technique (38):

- Wet hands with running water
- Place soap in palms
- Rub together to make a lather
- Scrub hands vigorously for 20 seconds
- Rinse soap off hands
- Dry hands with a disposable towel.
- Turn off the faucet by using a disposable towel

Handwashing with plain soap and running water mechanically removes soil and reduces the number of transient organisms on the skin, whereas antimicrobial soap kills or inhibits growth of both transient and resident flora (39, 40). All soaps also have the effect of dissolving the lipid envelope of enveloped viruses, and have cell wall effects that are bactericidal. Consideration for selection and use of soap products in practices include:

- Liquid or foam soap products should be selected rather than bar soaps to reduce the opportunity for cross-contamination
- Liquid soap dispensers should not be topped off. Refillable dispensers should be completely emptied, cleaned and then refilled to prevent creation of a bacterial reservoir
- Moisturizing soaps can preserve skin integrity and encourage compliance
- Plain or antibacterial products are appropriate for routine use

Alcohol-based gels are highly effective against bacteria and enveloped viruses and may be used if hands are not visibly soiled (35, 41, 42). Apply alcohol-based gel to palm of one hand and rub hands together until dry, covering all surfaces of hands and fingers. Follow the manufacturer's instructions regarding the amount of product to use (35). Alcohol-based gels are not effective against some non-enveloped viruses (e.g., norovirus, rotavirus, parvovirus), bacterial spores (e.g., anthrax, *Clostridium difficile*), or protozoal parasites (e.g., cryptosporidia) (35, 42).

Antimicrobial-impregnated wipes (i.e., towelettes), followed by alcohol-based gels, may be used when running water is not available. Used alone, wipes are not as effective as alcohol-based hand gels or washing hands with soap and running water (35).

## **2. Use of Gloves and Sleeves**

Gloves reduce the risk of pathogen transmission by providing barrier protection. They should be worn when touching blood, body fluids, secretions, excretions, mucous membranes, and non-intact skin. However, wearing gloves (including sleeves) does not replace handwashing (43, 44).

Gloves should be changed between examinations of individual animals or animal groups (e.g., litter of puppies/kittens, group of cattle) and between dirty and clean procedures on a single patient. Changing gloves between patients reduces the likelihood of spreading microorganisms from an animal or a fomite to another animal or person. Gloves should be worn when cleaning cages and environmental surfaces. They should be removed promptly after use, avoiding contact between skin and the outer glove surface. Disposable gloves should not be washed and reused (45, 46). Hands should be washed immediately after glove removal.

Gloves come in a variety of materials. Choice of gloves depends on their intended use. If latex allergies are a concern, acceptable alternatives include nitrile or vinyl gloves. For more information on preventing allergic reactions to natural rubber in the workplace, refer to the National Institute of Occupational Safety and Health (NIOSH) website (47).

Note: Gloves are not necessary when examining or handling normal, healthy animals.

### **3. Facial Protection**

Facial protection prevents exposure of mucous membranes of the eyes, nose and mouth to infectious materials. Facial protection should be used whenever exposures to splashes or sprays are likely to occur (4, 27).

Facial protection should include a mask worn with either goggles or a face shield. A surgical mask provides adequate protection during most veterinary procedures that generate potentially infectious aerosols. These include dentistry, nebulization, suctioning, bronchoscopy, lavage, flushing wounds and cleaning with high pressure sprayers.

### **4. Respiratory Protection**

Respiratory protection is designed to protect the respiratory tract from zoonotic infectious diseases transmitted through the air. The need for this type of protection is limited in veterinary medicine. However, it may be necessary in certain situations, such as when investigating abortion storms in small ruminants (Q fever), significant poultry mortality (avian influenza), ill psittacines (avian chlamydiosis) or other circumstances where there is concern about aerosol transmission. The N-95 rated disposable particulate respirator is a mask that is inexpensive, readily available, and easy to use. Fit-testing of the N-95 respirator is recommended but not required by the OSHA Respiratory Protection Standard (Appendix C). Respiratory evaluation and training should be provided for all personnel who use respirators (48). For additional information about fit-testing, contact the state labor department or visit OSHA's website at [www.osha.gov](http://www.osha.gov).

### **5. Protective Outerwear**

#### **a. Lab Coats, Smocks, and Coveralls**

This apparel is designed to protect street clothes or scrubs from contamination but is generally not fluid resistant, so they should not be used in situations where splashing or soaking with potentially infectious liquids is anticipated. For most personnel, this outerwear should be changed daily. Garments should be changed promptly whenever visibly soiled or contaminated with body fluids or feces. These garments should not be worn outside of the work environment (4, 49, 50).

#### **b. Non-Sterile Gowns**

Gowns provide more coverage for barrier protection than lab coats. Permeable gowns can be used for general care of patients in isolation. Impermeable gowns should be used to provide greater protection when splashes or large quantities of body fluids are present or anticipated. Disposable gowns should not be reused. Reusable fabric gowns should be laundered between each use. Gloves are indicated whenever gowns are worn. Gowns and gloves should be removed and placed in the trash or laundry bin before leaving the animal's environment. Hands should be washed immediately afterwards (27).

Employees should be trained to properly remove gowns to avoid contaminating themselves or their environments. The outer (contaminated) surface of a gown should only be touched with gloves.

- After unfastening ties, peel the gown from the shoulders and arms by pulling on the chest surface with gloves.

- Remove the gown, avoiding contact between the outer surface and clean surfaces.
- Ball the gown up for disposal while keeping the contaminated surface on the inside.
- Remove gloves and wash hands.
- If body fluids have soaked through the gown, promptly remove the contaminated clothing and wash the skin.

### **c. Footwear**

Footwear should be suitable for the specific working conditions (e.g., rubber boots for farm work) and should protect veterinary personnel from exposure to infectious material as well as trauma. Recommendations include shoes or boots with thick soles and closed-toe construction, which are impermeable to liquid, and easily cleaned. Disposable shoe covers or booties add an extra level of protection when heavy quantities of infectious materials are present or expected.

### **d. Headcovers**

Disposable headcovers provide a barrier when gross contamination of the hair and scalp is expected. Disposable headcovers should not be reused.

NOTE: The personal protective measures in this document do not eliminate the need for appropriate environmental engineering controls, such as hazard removal and safety engineering (e.g., separation of patient areas from break rooms). Personal protective equipment (e.g., respirators) should always be the last line of defense and should always be accompanied by appropriate training for personnel. While engineering control measures are beyond the scope of this document, more information is available at [www.osha.gov](http://www.osha.gov).

## **6. Bite and Other Animal-Related Injury Prevention**

During their careers, the majority (61%-68%) of veterinarians suffer an animal-related injury resulting in hospitalization and/or significant lost work time (1, 51-54). These are mainly dog and cat bites, kicks, cat scratches and crush injuries, and account for most occupational injuries among veterinarians (1, 51-54). In a recent study seeking to identify factors associated with increased risk of being bitten by a dog or cat in a veterinary teaching hospital, pets identified with a warning sign or considered more difficult to handle were four to five times more likely than other animals to have bitten a staff member while hospitalized. Yet only 47% of dogs and cats considered likely to bite were muzzled (compared to 12% to 14% of animals considered unlikely to bite) (55). Veterinary personnel reliably interpret the behaviors associated with an animal's propensity to bite; their professional judgment should be relied upon to guide bite prevention practices.

Approximately 3% to 18% of dog bites and 28% to 80% of cat bites become infected (56). Most clinically infected dog and cat bite wounds are mixed infections of aerobic and anaerobic bacteria. The most commonly isolated aerobes are *Pasteurella multocida* (cats), *Pasteurella canis* (dogs), streptococci, staphylococci, *Moraxella*, and *Neisseria weaverii*; the most commonly isolated anaerobes include *Fusobacterium*, *Bacteroides*,

*Porphyromonas*, and *Prevotella* (56). In addition, rare but serious systemic infections with invasive pathogens including *Capnocytophaga canimorsus*, *Bergeyella zoohelicum*, *Bartonella henselae*, and CDC nonoxidizer 1 group may occur following bites or scratches (56-60).

Veterinary personnel should take all necessary precautions to prevent animal-related injuries in the clinic and in the field. These may include physical restraints, bite-resistant gloves, muzzles, sedation, or anesthesia, and relying on experienced veterinary personnel rather than owners to restrain animals. Practitioners should remain alert for changes in their patients' behavior.

Veterinary personnel attending large animals should have an escape route in mind at all times (1, 54). When bites and scratches occur, immediate and thorough washing of the wound with soap and water is critical. Prompt medical attention should be sought for puncture wounds and other serious injuries. The need for tetanus immunization, antibiotics or rabies post-exposure prophylaxis should be evaluated. Animal bites may need to be reported to local or state public health agencies. Emergency contact information should be posted in the clinic.

## **B. Protective Actions During Veterinary Procedures**

### **1. Intake**

Waiting rooms should be a safe environment for clients, animals and employees. Aggressive or potentially infectious animals should be placed directly into an exam room. Animals with respiratory or gastrointestinal signs, or a history of exposure to a known infectious disease should be asked to enter through an alternative entrance to avoid traversing the reception area (61). If they must come through the reception area, they should be carried or placed on a gurney and taken directly into an exam room.

### **2. Examination of Animals**

All veterinary personnel must wash their hands between examinations of individual animals or animal groups (e.g., litter of puppies/kittens, herd of cattle). Hand hygiene is the most important measure to prevent transmission of zoonotic diseases while examining animals. Every exam room should have a sink with running water, a liquid soap dispenser, and paper towels. Alcohol-based hand gels may also be provided for use in conjunction with handwashing.

Veterinary personnel should wear protective outerwear and use gloves and other protective equipment appropriate for the situation. Potentially infectious animals should be examined in a dedicated exam room and should remain there until initial diagnostic procedures and treatments have been performed.

### **3. Injections, Venipuncture, and Aspirations**

#### **a. Needlestick Injury Prevention**

Needlestick injuries are among the most prevalent accidents in the veterinary workplace (62, 63). The most common needlestick injury is inadvertent injection of a vaccine (1, 64,

65). In a 1995 survey of 701 North Carolina veterinarians, 27% of respondents had accidentally self-inoculated rabies vaccine and 7% (23% of large animal veterinarians) live *Brucella* vaccine (1). Needle punctures sustained during procedures such as fine-needle aspiration are potential sources of zoonotic pathogens (66).

The most important precaution is to avoid recapping needles. Recapping causes more injuries than it prevents (67). When it is absolutely necessary to recap needles as part of a medical procedure or protocol, or if a puncture-proof, leak-resistant sharps container is not available, a mechanical device such as forceps can be used to replace the cap on the needle or the one-handed "scoop" technique may be employed (68). This technique involves holding the syringe with the attached needle or the needle hub alone (when unattached) and scooping or sliding the cap, which is lying on a horizontal surface, onto the needle's sharp end. Once the point of the needle is covered, the cap is tightened by pushing it against an object, or by pulling the base of the needle cap onto the hub of the needle with the same hand holding the syringe.

When injecting live vaccines or aspirating body fluids or tissue, the used syringe with the needle attached should be placed in a sharps container. Following most other veterinary procedures, the needle and syringe may be separated for disposal of the needle in the sharps container. This can be most safely accomplished by using the needle removal device on the sharps container, which allows the needle to drop directly into the container. Needles should never be removed from the syringe by hand. In addition, needle caps should not be removed by mouth.

Sharps containers are safe and economical, and should be located in every area where animal care occurs (69-71). Sharps should not be transferred from one container to another. Devices that cut needles prior to disposal should not be used because they increase the potential for aerosolization of the contents (69).

#### **b. Barrier Protection**

Gloves should be worn during venipuncture on animals suspected of having an infectious disease and when performing soft tissue aspirations. Currently, there is no data indicating that venipuncture on healthy animals carries a significant risk of infection.

#### **4. Dental Procedures**

Dental procedures create infectious aerosols and there is risk of exposure to splashes or sprays of saliva, blood, and infectious particles. There is also the potential for cuts and abrasions from dental equipment or teeth (72). The veterinary staff performing the dental procedure and anyone in the immediate vicinity (e.g. the veterinary anesthesiologist) should wear protective outerwear, gloves, mask, and a face shield or goggles. In one study, irrigating the oral cavity with a 0.12% chlorohexadine solution significantly decreased bacterial aerosolization (73).

#### **5. Resuscitations**

Resuscitations are particularly hazardous because they may occur without warning and unrecognized/undiagnosed zoonotic infectious agents may be involved. For example, a

dog that presents in respiratory failure after being hit by a car may have been in the road due to clinical rabies. Barrier precautions such as gloves, mask, and face shield or goggles should be worn at all times. Never blow into the nose/mouth of an animal or into an endotracheal tube to resuscitate an animal; instead, intubate the animal and use an ambubag or an anesthesia machine/respirator.

## **6. Obstetrics**

Common zoonotic agents, including *Brucella*, *Coxiella burnetii*, and *Listeria monocytogenes*, may be found in high concentrations in the birthing fluids of aborting or parturient animals, stillborn fetuses, and neonates (74). Gloves, sleeves, mask or respirator, face shield or goggles, and impermeable protective outerwear should be employed as needed to prevent exposures to potentially infectious materials. During resuscitation, do not blow into the nose or mouth of a non-respiring neonate.

## **7. Necropsy**

Necropsy is a high risk procedure due to contact with infectious body fluids, aerosols, and contaminated sharps. Non-essential persons should not be present. Veterinary personnel involved in or present at necropsies should wear gloves, masks, face shields or goggles and impermeable protective outerwear as needed. In addition, cut-proof gloves should be used to prevent sharps injuries. Respiratory protection (including environmental controls and respirators) should be employed when band saws or other power equipment are used.

Decisions regarding whether to perform necropsy on animals suspected of having a notifiable infectious disease or foreign animal disease should be made in consultation with the state veterinarian.

## **8. Diagnostic Specimen Handling**

Feces, urine, aspirates, and swabs should be presumed to be infectious. Protective outerwear and disposable gloves should be worn when handling these specimens. Discard gloves and wash hands before touching clean items (e.g., microscopes, telephones, food).

Although in veterinary practices animal blood specimens have not been a significant source of occupational infection, percutaneous and mucosal exposure to blood and blood products should be avoided.

Eating and drinking must not be allowed in the laboratory.

## **C. Environmental Infection Control**

### **1. Isolation of Infectious Animals**

Patients with a contagious or zoonotic disease should be clearly identified so their infection status is obvious to everyone, including visitors allowed access to clinical areas. Prominent signage should indicate that the animal may be infectious and should outline any additional precautions that should be taken (27, 75).

Ideally, veterinary practices should utilize a single-purpose isolation room for caring for and housing contagious patients (75). Alternatively, a dedicated exam room that can be easily emptied of non-essential equipment, cleaned and disinfected can be transformed into an isolation room. A mobile cage unit may be brought in for exclusive use by the infectious animal. If an isolation room has negative pressure air handling, air pressures should be monitored daily while in use and the air should be exhausted outside of the building, away from animal and public access areas, employee break areas, and air intake vents (4, 75, 76).

Only the equipment and materials needed for the care and treatment of the patient should be kept in the isolation room. Items intended for use in the isolation room should remain in this area and duplicate new items purchased for use elsewhere in the hospital. When necessary, items removed from the isolation area should be taken apart, cleaned, and disinfected prior to removal. Use of disposable articles can minimize the need to bring soiled items out of the isolation room. Access to the isolation room should be limited and a sign-in sheet should be kept of all people having contact with a patient in isolation (75, 77).

Limited data are available on the efficacy of footbaths. When used, a disinfectant footbath should be placed just inside the door of the isolation area and used before departing the room (75, 78). Footbath disinfectant should be changed daily or when visibly dirty. If shoe or boot coverings are used, personnel should be trained to use, remove, and dispose of them properly.

Depending on the diagnosis and the mode of transmission of the disease, clean (non-sterile) gowns, shoe covers, gloves, masks and eye protection should be worn when handling an animal with a zoonotic disease. The gloves should be discarded, but typically the rest of the personal protective equipment (e.g., gown, mask) may be re-used and should remain in the isolation room with the patient. However, if the gown, mask, or other protective equipment is contaminated with body fluids, it should be replaced. Protective equipment should be cleaned and disinfected between patients. Potentially contaminated materials should be bagged before transport within the practice and disinfected or disposed of appropriately according to their level of hazard. In many cases, all the materials used in the isolation room would be treated as bio-waste (27, 71, 75).

## **2. Cleaning and Disinfection\* of Equipment and Environmental Surfaces**

Proper cleaning of environmental surfaces, including work areas and equipment, prevents transmission of zoonotic pathogens. Environmental surfaces and equipment should be cleaned between uses or whenever visibly soiled. A recent report indicates that directed misting application of a peroxygen disinfectant for environmental decontamination is effective in veterinary settings (79).

Surfaces where animals are housed, examined, or treated should be made of non-porous, easily cleanable materials. Surfaces should be cleaned to remove gross contamination before disinfection because organic material decreases the effectiveness of most disinfectants (80). When cleaning, avoid generating dust that may contain pathogens by



using central vacuum units, wet mopping, dust mopping, or electrostatic sweeping (e.g., Swiffer®). Surfaces may be lightly sprayed with water prior to mopping or sweeping. Areas to be cleaned should be appropriately ventilated.

Clean items should be kept separate from dirty items. Gloves should be worn when cleaning equipment, animal cages, and surfaces. Clean and disinfect equipment according to its intended use, the manufacturer's recommendations, and practice policy. Equipment must be cleaned before sterilization or chemical disinfection. Exposure to aerosols generated by brushes during cleaning can be minimized by implementing preventive work practices, such as wearing facial protection and containing splatter.

Normal dishwashing of food and water bowls is adequate for hospitalized patients with infectious diseases (4), although disposable dishes might be considered for animals hospitalized in isolation. Toys, litter boxes, and other miscellaneous items should be discarded or cleaned and disinfected between patients. If these items are visibly soiled, gloves should be worn. Litter boxes should be cleaned or disposed of at least daily by a non-pregnant staff member.

Hands should be washed after finishing a cleaning activity.

\*NOTE: Environmental Protection Agency (EPA)-registered hospital disinfectants (see Appendix B) should be used. To ensure effectiveness, disinfectants should be used according to manufacturers' instructions, with particular regard to proper dilution and contact time. Personnel engaged in cleaning should be trained in safe practices and should be provided necessary safety equipment according to the product's Material Safety Data Sheet (MSDS).

### **3. Handling of Laundry**

Although soiled laundry may be contaminated with pathogenic microorganisms, the risk of disease transmission is negligible if handled correctly. Gloves should always be worn when handling soiled laundry. Bedding and other laundry should be machine washed with standard laundry detergent and machine dried. To prevent cross-contamination, separate storage and transport bins should be used for clean and dirty laundry.

### **4. Decontamination and Spill Response**

Spills and splashes of blood or other body fluids should be immediately sprayed with disinfectant and contained by dropping absorbent material (e.g., paper towels, sawdust, cat litter) on them. A staff person should wear gloves, a mask, and protective clothing (including shoe covers if the spill is on the floor and may be stepped in) before beginning the clean-up. The spilled material should be picked up and sealed in leak-proof plastic bags. After the spilled material is removed, the area should be cleaned, and disinfected according to the manufacturer's instructions. Clients, patients, and employees not involved in the clean-up should be kept away from the area until disinfection is completed (76).

## **5. Veterinary Medical Waste**

Veterinary medical waste is a potential source of zoonotic pathogens if not handled appropriately (71, 81). Medical waste is defined and regulated at the state level by multiple agencies, but may include sharps, tissues, contaminated materials, and dead animals. AVMA recommends voluntary compliance with the OSHA Bloodborne Pathogen Standard (82) in order to best protect veterinary personnel. It is beyond the scope of this Compendium to describe veterinary medical waste management in detail. Consult with local and/or state health departments and municipal governments for guidance. Additional information and links to state regulating agencies can be found on the Environmental Protection Agency website: [www.epa.gov/epaoswer/other/medical](http://www.epa.gov/epaoswer/other/medical).

## **6. Rodent and Vector Control**

Many important zoonotic pathogens are transmitted by rodents or insect vectors. The principles of integrated pest management (IPM) are central to effective prevention and control (83, 84). IPM practices include:

- Sealing entry and exit points into buildings. Common methods include the use of caulk, steel wool, or lath metal under doors and around pipes (85)
- Storing food and garbage in metal or thick plastic containers with tight lids
- Disposing of food-waste promptly
- Eliminating potential rodent nesting sites (e.g., clutter, hay storage)
- Maintaining snap traps throughout the practice to trap rodents (check daily)
- Removing sources of standing water (empty cans, tires, etc.) from around the building to prevent breeding of mosquitoes (86)
- Installing and maintaining window screens to prevent entry of insects into buildings (86)

Additional measures may be warranted for control of specific pests. Veterinary practices may wish to contact a pest control company for additional guidance.

## **7. Other Environmental Controls**

Designated staff areas should be set aside for eating, drinking, smoking, etc. These activities should never occur in patient care areas. Separate refrigerators should be used for human food, animal food, and biologics. Dishes for human use should be cleaned and stored away from animal care areas.

# **IV. EMPLOYEE HEALTH**

## **A. General**

Veterinary practice managers should promote infection control as part of a comprehensive program of employee health. Senior management support for workplace safety programs has been significantly associated with the level of staff compliance with safety policies and procedures (87).

## **1. Infection Control Personnel**

Veterinary practices should designate a staff member to be responsible for developing and maintaining infection control policies, keeping records and managing workplace exposure and injury incidents.

## **2. Employee Immunization Policies and Record Keeping**

### **a. Rabies**

Veterinary personnel who have contact with animals should be vaccinated against rabies in accordance with recommendations of the Advisory Committee on Immunization Practices (ACIP) (88). Preexposure rabies vaccination consists of three doses of a licensed human rabies vaccine administered on days 0, 7, and 21 or 28. In addition to preexposure rabies vaccination, the ACIP guidelines recommend a rabies antibody titer be checked every two years for those in the frequent risk category, including veterinarians and their animal handling staff. Preexposure vaccination for rabies does not eliminate the need for appropriate treatment following a known rabies exposure, but it does simplify the postexposure treatment regimen (2 doses of vaccine without rabies immune globulin instead of rabies immune globulin plus 5 doses of vaccine). In addition, preexposure vaccination may protect against unrecognized rabies exposures or when postexposure treatment is delayed (88).

### **b. Tetanus**

All staff should have an initial series of tetanus immunizations, followed by a booster vaccination every 10 years. In the event of a possible exposure to tetanus, such as a puncture wound, employees should be evaluated by their health care provider; a tetanus booster may be indicated (89).

### **c. Seasonal Influenza**

Veterinary personnel are encouraged to receive the current seasonal influenza vaccine, unless contraindicated. This is intended to minimize the small possibility that dual infection of an individual with human and avian or swine influenza virus could result in a new hybrid strain of the virus. *Check with CDC for updated recommendations* (90).

### **d. Immunization and Other Health Records**

Veterinary practices should maintain records on immunizations, rabies titers, exposures and emergency contact information for staff. This allows for an efficient response to occupational health incidents, such as rabies exposures, by providing necessary records to healthcare providers (91, 92). Records should be maintained in a retrievable, secure database (91). Maintaining these records will facilitate monitoring the work-related health status of employees (e.g., identifying when an employee is due for a rabies titer). Employee health records should be collected on a voluntary basis, with a clear understanding that confidentiality will be maintained. Other health-related issues that may influence employees' work duties should be documented in personnel files. Employees should inform their supervisor of changes in health status (e.g. pregnancy) that may affect work duties. All employees should inform their personal physicians that their work duties involve animal contact.

### **3. Staff Training and Education**

Staff training and education are essential components of an effective employee health program. All employees should receive education and training on injury prevention and infection control at the beginning of their employment and at least annually. Additional in-service training should be provided as recommendations change or if problems with infection control policies are identified. Training should emphasize the potential for zoonotic disease exposure and hazards associated with work duties, and include animal handling, restraint, and behavioral cue recognition. Staff participation in training should be documented.

### **4. Documenting Exposure Incidents**

Injuries or potential exposures to zoonotic pathogens should be reported to the practice owner or designee. Practice managers should develop policies that encourage reporting (92).

Any incident involving injury or a zoonotic pathogen should be thoroughly investigated and documented. Specific risk situations such as an immunocompromising condition or pregnancy and immunization status of those involved in the incident should be reviewed (92). A data collection form (e.g. OSHA form 300) should include:

- Time, date and location of the incident
- Person(s) involved
- Nature of the incident (injury, exposure or both) and a detailed description of the circumstances.
- Status of the animal involved (vaccination status, clinical condition, any infectious disease diagnostic test results or tests pending)
- Plans for follow up (evaluation by physician, reporting to public health)

Practices should contact their local or state health department to inquire about required reporting of bite incidents and zoonotic diseases.

## **B. Immunocompromised Personnel**

Immune deficiencies may put veterinary personnel at increased risk for acquiring zoonotic infections. Additionally, immunocompromised personnel are more likely to develop serious complications from infections. Immune deficiencies may result from underlying medical conditions (e.g., HIV/AIDS, diabetes mellitus, asplenia, pregnancy, certain malignancies), therapy for a variety of conditions (e.g., steroids, chemotherapeutic and immunosuppressive agents, radiation) or may be congenital. Immunocompromised employees and their supervisors should be aware of the following workplace encounters that may result in exposure to zoonotic pathogens:

- Processing laboratory samples
- Direct patient care, especially with the following high risk animals (38):
  - Young animals (ruminants prior to weaning, dogs and cats less than six months of age)
  - Animals with diarrhea
  - Parturient animals

- Stray or feral animals (especially predators of rodents and wildlife)
- Animals fed raw meat diets
- Reptiles or exotic, imported species
- Animals housed in crowded conditions (e.g., shelters)
- Unvaccinated animals or those with untreated internal or external parasites.

Data are limited regarding the risks of zoonotic infection for HIV-infected persons employed in veterinary settings and none exist to justify their exclusion. The risks associated with exposure to zoonotic pathogens in the workplace can be mitigated by appropriate infection control measures (93).

During pregnancy, women experience physiologic suppression of cell-mediated immunity, increasing their susceptibility to certain infections. These include toxoplasmosis, lymphocytic choriomeningitis virus infection, brucellosis, listeriosis and *Chlamydia psittaci*. Vertical transmission of certain zoonoses may result in abortion, stillbirth, prematurity or congenital anomalies.

Employees with immune dysfunction should discuss their status with the practice manager or owner so the practice can provide appropriate workplace accommodations to protect them. The use of infection control measures and personal protective equipment will reduce the risk of infection. In some cases, it may be advisable to consult the employee's healthcare provider or an infection control, public health or occupational health specialist in managing the zoonotic disease risk (94). Employers must abide by state and federal laws that protect pregnant women and persons with disabilities. The employee should be assured that confidential information will not be disclosed to others.

## **V. CREATING A WRITTEN INFECTION CONTROL PLAN**

All veterinary practices should have a written infection control plan, which should be reviewed and updated at least annually. Appendix D provides a model plan that can be tailored to individual practice needs. A modifiable electronic version is available on the NASPHV website at [www.nasphv.org](http://www.nasphv.org).

Effective infection control plans should:

- Be specific to the facility and practice type
- Be flexible to easily address new issues and incorporate new knowledge
- Provide explicit, well organized, understandable guidance
- Clearly describe the role of each staff member
- Be incorporated into new employee training and regularly reviewed with staff
- Include a process for the evaluation of infection control practices
- Be kept in work areas for quick reference
- Provide contact information, resources, and references (e.g., reportable disease list, public health contacts, local rabies codes and environmental health regulations, OSHA requirements, websites of interest, client education materials)

## **A. Communicating and Updating the Infection Control Plan**

**1. Availability:** Keep copies of the infection control plan and resource documents at locations readily accessible to all staff including reception, administrative, animal care, housekeeping and veterinary personnel.

**2. Leadership:** Senior and managerial personnel should set the standard for infection control practices, stress its importance to other staff and reference the infection control plan in daily activities.

**3. New Staff:** New staff should be given a copy of the infection control plan and receive detailed training on the practice's infection control procedures, staff vaccination recommendations, and how to report exposure incidents. Some employers may ask new staff members to sign a form stating they have received and read the plan.

**4. Continuing Education:** Infection control procedures should be reviewed regularly with staff at staff meetings, and veterinary Continuing Education on zoonotic diseases should be encouraged.

**5. Review and Revision:** A designated staff person should be responsible for regularly reviewing and revising the infection control plan as needed when new information becomes available or when clinical practices change. When revisions are made, they should be shared with all staff members and all copies of the plan updated at the same time.

**6. Assurance:** A designated staff person should be responsible for assuring the plan components are being carried out consistently and correctly. This person should also ensure that staff are counseled and/or corrective measures are instituted when deficiencies in infection control procedures are identified.

## VI. REFERENCES

1. Langley RL, Pryor WH, O'Brien KF. Health hazards among veterinarians: a survey and review of the literature. *J Agromedicine* 1995;2:23-52.
2. Snow J, Rice J. Infection control in veterinary clinics. *Northwest Public Health* 2005; Fall/Winter:22-23.
3. Jung S. Analysis of infection control practices among small animal veterinarians. MPH thesis, Rollins School of Public Health, Emory University 2006.
4. Garner JS. Guideline for isolation precautions in hospitals. The Hospital Infection Control Practices Advisory Committee. *Infect Control Hosp Epidemiol* 1996;17:53-80.
5. Marano N, Pappaioanou M. Historical, new, and reemerging links between human and animal health. *Emerg Infect Dis* 2004;10:2065-2066.
6. Taylor LH, Latham SM, Woolhouse ME. Risk factors for human disease emergence. *Philos Trans R Soc Lond B Biol Sci* 2001;356:983-989.
7. CDC. Update: multistate outbreak of monkeypox - Illinois, Indiana, Kansas, Missouri, Ohio, and Wisconsin, 2003. *Morb Mortal Wkly Rep* 2003; 52:642-646.
8. Wright, JG, Tengelsen LA, Smith KE, et al. Multi-drug resistant *Salmonella* Typhimurium in four animal facilities. *Emerg Infect Dis* 2005;11:1235-1241.
9. Cherry B, Burns A, Johnson GS, et al. *Salmonella* Typhimurium outbreak associated with veterinary clinic. *Emerg Infect Dis* 2004;10:2249-2251.
10. Bender JB, Tsukayama DT. Horses and the risk of zoonotic infections. *Vet Clin North Am Equine Pract* 2004;20:643-653.
11. Levine JF, Levy MG, Walker RL, et al. Cryptosporidiosis in veterinary students. *J Am Vet Med Assoc* 1988;193:1413-1414.
12. Anderson BC, Donndelinger T, Wilkins RM, et al. Cryptosporidiosis in a veterinary student. *J Am Vet Med Assoc* 1982;180:408-409.
13. Reif JS, Wimmer L, Smith JA, et al. Human cryptosporidiosis associated with an epizootic in calves. *Am J Public Health* 1989;79:1528-1530.
14. Pohjola S, Oksanen H, Jokipii L, et al. Outbreak of cryptosporidiosis among veterinary students. *Scand J Infect Dis* 1986;18:173-178.
15. Preiser G, Preiser L, Madeo L. An outbreak of cryptosporidiosis among veterinary science students who work with calves. *J Am Coll Health* 2003;51:213-215.
16. Gage KL, Dennis DT, Orloski KA, et al. Cases of cat-associated human plague in the Western US, 1977-1998. *Clin Infect Dis* 2000;30:893-900.
17. Dunstan RW, Reimann KA, Langham RF. Feline sporotrichosis. In: *Zoonosis Updates from the Journal of the American Veterinary Medical Association*. 2<sup>nd</sup> ed. Schaumburg: American Veterinary Medical Association, 1995;79-82.
18. Dunstan RW, Langham RF, Reimann KA, et al. Feline sporotrichosis: a report of five cases with transmission to humans. *J Am Acad Dermatol* 1986;15:37-45.
19. Nusbaum BP, Gulbas N, Horwitz SN. Sporotrichosis acquired from a cat. *J Am Acad Dermatol* 1983;8:386-391.
20. Reed KD, Moore FM, Geiger GE, et al. Zoonotic transmission of sporotrichosis: case report and review. *Clin Infect Dis* 1993;16:384-387.
21. Clinkenbeard KD. Diagnostic cytology: sporotrichosis. *Compend Contin Educ Pract Vet* 1991;13:207-211.
22. Baptiste KE, Williams K, Williams NJ, et al. Methicillin-resistant staphylococci in companion animals. *Emerg Infect Dis* 2005; 11:1942-1944.
23. Bender JB, Torres SMF, Gilbert SM, et al. Isolation of methicillin-resistant *Staphylococcus aureus* from a non-healing abscess in a cat. *Vet Rec* 2005;157:388-389.
24. Weese JS, Dick H, Willey BM, et al. Suspected transmission of methicillin-resistant *Staphylococcus aureus* between domestic pets and humans in veterinary clinics and in the household. *Vet Microbiol* 2006;115:148-155.
25. O'Mahony R, Abbott Y, Leonard FC, et al. Methicillin-resistant *Staphylococcus aureus* (MRSA) isolated from animals and veterinary personnel in Ireland. *Vet Microbiol* 2005;109:285-296.
26. Constable PJ, Harrington JM. Risks of zoonoses in a veterinary service. *BMJ* 1982;284:246-248.
27. Weese JS. Barrier precautions, isolation protocols, and personal hygiene in veterinary hospitals. *Vet Clin North Amer Equine Pract* 2004;20:543-559.

28. Hannah, Harold W. A veterinarian's liability to employees. *J Am Vet Med Assoc* 1994;204:361-362.
29. Smith R, Stiltz P. Controlling workers' compensation losses. *J Am Vet Med Assoc* 1996;209:526.
30. Lenhart SW, Seitz T, Trout D, et al. Issues affecting respirator selection for workers exposed to infectious aerosols: emphasis on healthcare settings. *Appl Biosafety* 2004;9:20-36.
31. Acha PN, Szyfres B. *Zoonoses and Communicable Diseases Common to Man and Animals. Vol 2: Chlamydioses, Rickettsioses, and Viroses.* 3<sup>rd</sup> ed. Washington D.C.: Pan American Health Organization, 2003;22.
32. Marrie TJ. Q Fever. In: Palmer SR, Soulsby L, Simpson DIH, eds. *Zoonoses*. New York: Oxford University Press, Inc, 1998;171-185.
33. McQuiston JH, Childs JE. Q fever in humans and animals in the United States. *Vector Borne Zoonotic Dis* 2002;2:179-191.
34. Nation PN, Fanning EA, Hopf HB, et al. Observations on animal and human health during the outbreak of *Mycobacterium bovis* in game farm wapiti in Alberta. *Can Vet J* 1999;40:113-117.
35. Boyce JM, Pittet D. Guideline for hand hygiene in health-care settings. *Morb Mortal Wkly Rep Recomm Rep* 2002;51:1-48.
36. Larson EL. APIC guideline for handwashing and hand antisepsis in health care settings. *Am J Infect Control* 1995;23:251-269.
37. Lin CM, Wu FM, Kim HK, et al. A comparison of hand washing techniques to remove *Escherichia coli* and caliciviruses under natural or artificial nails. *J Food Protect* 2003;66:2296-2301.
38. National Association of State Public Health Veterinarians. Compendium of measures to prevent disease associated with animals in public settings, 2006. Available at: <http://s94745432.onlinehome.us/AnimalsInPublicSettings.pdf>. Accessed Aug 22, 2006.
39. Ayliffe, GA, Babb JR, Quoraishi AH. A test for hygienic hand disinfection. *J Clin Pathol* 1978;31:923-928.
40. Steere AC, Mallison GF. Handwashing practices for the prevention of nosocomial infections. *Ann Intern Med* 1975;83:683-690.
41. Widmer AF, Dangel M. Alcohol-based handrub: evaluation of technique and microbiological efficacy with international infection control professionals. *Infect Control Hosp Epidemiol* 2004;25:207-209.
42. Sickbert-Bennett EE, Weber DJ, Gergen-Teague MF, et al. Comparative efficacy of hand hygiene agents in the reduction of bacteria and viruses. *Am J Infect Control* 2005;33:67-77.
43. Goldmann DA. The role of barrier precautions in infection control. *J Hosp Infect* 1991;18:515-523.
44. Olsen RJ, Lynch P, Coyle MB, et al. Examination gloves as barriers to hand contamination in clinical practice. *J Amer Med Assoc* 1993;2:350-353.
45. Doebbeling BN, Pfaller MA, Houston AK, et al. Removal of nosocomial pathogens from the contaminated glove: implications for glove reuse and handwashing. *Ann Intern Med* 1988;109:394-398.
46. Patterson JE, Vecchio J, Pantelick EL, et al. Association of contaminated gloves with transmission of *Acinetobacter calcoaceticus* var. *anitratus* in an intensive care unit. *Am J Med* 1991;91:479-483.
47. National Institute for Occupational Safety and Health (NIOSH). Preventing allergic reactions to natural rubber latex in the workplace. DHHS (NIOSH) Publication No. 97-135. Available at: [www.cdc.gov/niosh/latexalt.html](http://www.cdc.gov/niosh/latexalt.html). Accessed Aug 22, 2006.
48. U.S. Department of Labor Occupational Safety and Health Administration. Occupational safety and health standards: personal protective equipment. 1910.134, Appendix A: Fit testing procedures (mandatory).
49. Belkin NL. Use of scrubs and related apparel in health care facilities. *Am J Infect Control* 1997;5:401-404.
50. Belkin NL. Home laundering of soiled surgical scrubs: surgical site infections and the home environment. *Am J Infect Control* 2001;29:58-64.
51. Landercasper J, Cogbill TH, Strutt PJ, et al. Trauma and the veterinarian. *J Trauma* 1988;28:1255-1259.
52. Gabel CL, Gerberich SG. Risk factors for injury among veterinarians. *Epidemiology* 2002;13:80-86.
53. Jeyaretnam J, Jones H, Phillips M. Disease and injury among veterinarians. *Aust Vet J* 2000;78:625-629.
54. Neinhaus A, Skudlik C, Seidler A. Work-related accidents and occupational diseases in veterinarians and their staff. *Int Arch Occup Environ Health* 2005;78:230-238.
55. Drobatz KJ, Smith G. Evaluation of risk factors for bite wounds inflicted on caregivers by dogs and cats in a veterinary teaching hospital. *J Am Vet Med Assoc* 2003;223:312-316.
56. Talan DA, Citron DM, Abrahamian FM, et al. Bacteriologic analysis of infected dog and cat bites. *N Engl J Med* 1999;340:85-92.
57. Le Moal G, Landron C, Grollier G, et al. Meningitis due to *Capnocytophaga canimorsus* after receipt of a dog bite: case report and review of the literature. *Clin Infect Dis* 2003;36:e42-e46.



58. Shukla SK, Paustian DL, Stockwell PJ, et al. Isolation of a fastidious *Bergeyella* species associated with cellulitis after a cat bite and a phylogenetic comparison with *Bergeyella zoohelcum* strains. *J Clin Microbiol* 2004;42:290-293.
59. Kaiser RM, Garman RL, Bruce MG, et al. Clinical significance and epidemiology of NO-1, an unusual bacterium associated with dog and cat bites. *Emerg Infect Dis* 2002;8:171-174.
60. Hara H, Ochiai T, Morishima T, et al. *Pasteurella canis* osteomyelitis and cutaneous abscess after a domestic dog bite. *J Am Acad Dermatol* 2002;s151-s152.
61. CDC. Guidelines and resources, Monkeypox infections in animals: updated interim guidance for veterinarians. Available at: [www.cdc.gov/ncidod/monkeypox/animalguidance.htm](http://www.cdc.gov/ncidod/monkeypox/animalguidance.htm). Accessed Aug 25, 2006.
62. Poole AG, Shane SM, Kearney MT, et al. Survey of occupational hazards in companion animal practices. *J Am Vet Med Assoc* 1998;212:1386-1388.
63. Poole AG, Shane SM, Kearney MT, et al. Survey of occupational hazards in large animal practices. *J Am Vet Med Assoc* 1999;215:1433-1435.
64. Hafer AL, Langley RL, Morrow WEM, et al. Occupational hazards reported by swine veterinarians in the United States. *J Swine Health Prod* 1996;4:128-141.
65. Wilkins JR, Bowman ME. Needlestick injuries among female veterinarians: frequency, syringe contents and side-effects. *Occup Med* 1997;47:451-457.
66. Ramsey DT. Blastomycosis in a veterinarian. *J Am Vet Med Assoc* 1994;205:968.
67. U.S. Department of Labor Occupational Safety & Health Administration. Bloodborne and needlestick prevention: possible solutions. Available at: [www.osha.gov/SLTC/bloodbornepathogens/solutions.html](http://www.osha.gov/SLTC/bloodbornepathogens/solutions.html). Accessed Aug 22, 2006.
68. Cornell Center for Animal Resources and Education. CARE711.01 Sharps precautions. Available at: [www.research.cornell.edu/care/CARE711.pdf](http://www.research.cornell.edu/care/CARE711.pdf). Accessed Aug 22, 2006.
69. Seibert PJ. Hazards in the hospital. *J Am Vet Med Assoc* 1994;204:352-360.
70. Grizzle WE, Fredenburgh J. Avoiding biohazards in medical, veterinary, and research laboratories. *Biotech Histochem* 2001;76:183-206.
71. Brody MD. Safety in the veterinary medical workplace environment. *Vet Clin North Am Small Anim Pract* 1993;23:1071-1084.
72. Holmstrom SE, Bellows J, Colmery B, et al. AAHA dental care guidelines for dogs and cats. *J Am Anim Hosp Assoc* 2005;41:277-283.
73. Logothetis DD, Martinez-Welles JM. Reducing bacterial aerosol contamination with a chlorhexidine gluconate pre-rinse. *J Am Dent Assoc* 1995;126:1634-1639.
74. Heymann DL, ed. Control of Communicable Diseases Manual. 18<sup>th</sup> ed. Washington D.C.: American Public Health Association, 2004;75-78,434-438,309-312.
75. Contagious disease standards – effective containment. American Animal Hospital Association 2005.
76. CDC. Guidelines for environmental infection control in health-care facilities: recommendations of CDC and the Healthcare Infection Control Practices Advisory Committee (HICPAC). *Morb Mortal Wkly Rep Recomm Rep* 2003;52:1-42.
77. Virginia-Maryland Regional College of Veterinary Medicine. Policy Number 9210 – Isolation and infection control policy and procedures. Available at: [www.vth.vt.edu/policies/showpnp.asp?id=62](http://www.vth.vt.edu/policies/showpnp.asp?id=62). Accessed Aug 22, 2006.
78. Morley PS, Morris SN, Hyatt DR, et al. Evaluation of the efficacy of disinfectant footbaths as used in veterinary hospitals. *J Am Vet Med Assoc* 2005;226:2053-2058.
79. Patterson G, Morley PS, Blehm KD, et al. Efficacy of directed misting application of a peroxygen disinfectant for environmental decontamination of a veterinary hospital. *J Am Vet Med Assoc* 2005;227:597-602.
80. Dwyer RM. Environmental disinfection to control equine infectious diseases. *Vet Clin North Amer Equine Pract* 2004;20:531-542.
81. Brody MD. AVMA guide for veterinary medical waste management. *J Am Vet Med Assoc* 1989;195:440-452.
82. U.S. Department of Labor Occupational Safety and Health Administration. Occupational safety and health standards: toxic and hazardous substances. 1910.1030: Bloodborne pathogens.
83. Kogan M. Integrated pest management: historical perspectives and contemporary developments. *Annu Rev Entomol* 1998;43:243-270.

84. Peter RJ, Van den Bossche P, Penzhorn BL, et al. Tick, fly, and mosquito control—lessons from the past, solutions for the future. *Vet Parasitol* 2005;132:205-215.
85. CDC. Hantavirus pulmonary syndrome. Available at: [www.cdc.gov/ncidod/diseases/hanta/hps\\_stc/stc\\_seal.htm](http://www.cdc.gov/ncidod/diseases/hanta/hps_stc/stc_seal.htm). Accessed Aug 22, 2006.
86. CDC. West Nile Virus. Available at: [www.cdc.gov/ncidod/dvbid/westnile/prevention\\_info.htm](http://www.cdc.gov/ncidod/dvbid/westnile/prevention_info.htm). Accessed Aug 22, 2006.
87. Gershon RRM, Karkashian CD, Grosch JW, et al. Hospital safety climate and its relationship with safe work practices and workplace exposure incidents. *Am J Infect Control* 2000;28:211-221.
88. CDC. Human rabies prevention — United States, 1999: Recommendations of the Advisory Committee on Immunization Practices (ACIP). *Morb Mortal Wkly Rep Recomm Rep* 1999;48:1-21.
89. CDC. Immunization of health-care workers: Recommendations of the Advisory Committee on Immunization Practices (ACIP) and the Hospital Infection Control Practices Advisory Committee (HICPAC). *Morb Mortal Wkly Rep Recomm Rep* 1997;46:1-44.
90. CDC. Interim guidance for protection of persons involved in U.S. avian influenza outbreak disease control and eradication activities, 2006. Available at: [www.cdc.gov/flu/avian/professional/protect-guid.htm](http://www.cdc.gov/flu/avian/professional/protect-guid.htm). Accessed Aug 22, 2006.
91. Bolyard EA, Tablan OC, Williams WW, et al. Guideline for infection control in health care personnel, 1998. *Am J Infect Control* 1998;26:289-354.
92. Herwaldt LA, Pottinger JM, Carter CD, et al. Exposure workups. *Infect Control Hosp Epidemiol* 1997;18:850-871.
93. CDC. Guidelines for preventing opportunistic infections among HIV-infected persons—2002: Recommendations of the U.S. Public Health Service and the Infectious Diseases Society of America. *Morb Mortal Wkly Rep Recomm Rep* 2002;51:1-52.
94. Grant S, Olsen CW. Preventing zoonotic diseases in immunocompromised persons: the role of physicians and veterinarians. *Emerg Infect Dis* 1999;5:159-163.

## APPENDIX A

### Zoonotic Diseases of Importance in the United States

Disease	Agent	Means of transmission to humans	Most common species associated with transmission to humans	Nationally notifiable for humans (H) or animals (A)	Severe disease usually associated with immune suppression	Fatalities reported in humans
<b>Acariasis (Mange)</b>	<i>Sarcoptes scabiei</i> , <i>Notoedres cati</i> and other species of mites	contact	Dogs, cats, horses, goats, sheep, swine, birds			No
<b>Anthrax</b>	<i>Bacillus anthracis</i>	contact, aerosol, vector	cattle, sheep, goats, horses	H, A		Yes
<b>Avian Influenza</b>	<i>Highly Pathogenic AI viruses</i>	contact, aerosol	poultry, pet birds	H,A		Yes
<b>Babesiosis</b>	<i>Babesia microti</i> and other species	vector	cattle		✓	Yes
<b>Baylisascariasis</b>	<i>Baylisascaris procyonis</i>	contact	raccoons			Yes
<b><i>Bordetella bronchiseptica</i> infection</b>	<i>Bordetella bronchiseptica</i>	aerosol	dogs, pigs, rabbits, guinea pigs		✓	No
<b>Brucellosis</b>	<i>Brucella melitensis</i> , <i>B. abortus</i> , <i>B. suis</i> , <i>B. canis</i>	aerosol, contact	goats, cattle, swine, dogs, horses	H, A		Yes
<b>Campylobacteriosis</b>	<i>Campylobacter jejuni</i> , <i>C. fetus</i> , <i>C. coli</i>	contact	cattle, sheep, goats, pigs, dogs, cats, birds, mink, ferrets, hamsters			Rare
<b><i>Capnocytophaga canimorsus</i> infection</b>	<i>Capnocytophaga canimorsus</i> , <i>C. cynodegmi</i>	contact	dog, cat		✓	Yes
<b>Cat scratch disease</b>	<i>Bartonella henselae</i>	contact	cats		✓	Rare
<b>Chlamydiosis (mammalian)</b>	<i>Chlamydomydia abortus</i> , <i>C. felis</i>	aerosol, contact	cattle, sheep, goats, llamas, cats			Yes
<b>Contagious pustular dermatitis (orf, contagious ecthyma)</b>	Parapoxvirus	contact	sheep, goats			No
<b>Cryptococcosis</b>	<i>Cryptococcus neoformans</i>	aerosol	cats, pigeons, cockatoos		✓	Yes
<b>Cryptosporidiosis</b>	<i>Cryptosporidium parvum</i>	contact	cattle, typically calves	H	✓	Yes
<b>Dermatophilosis</b>	<i>Dermatophilus congolensis</i>	contact, vector	goats, sheep, cattle, horses			No
<b>Dermatophytosis (Ringworm)</b>	<i>Microsporum</i> sp., <i>Trichophyton</i> sp. <i>Epidermophyton</i> sp.	contact	cats, dogs, cattle, goats, sheep, horses, lagomorphs, rodents		✓	No
<b><i>Dipylidium</i> infection (Tapeworms)</b>	<i>Dipylidium caninum</i>	vector	dog, cat			No
<b><i>E. coli</i> O157:H7 infection</b>	<i>Escherichia coli</i> O157:H7	contact	cattle, goats, sheep	H		Yes

## Zoonotic Diseases of Importance in the United States

Disease	Agent	Means of transmission to humans	Most common species associated with transmission to humans	Nationally notifiable for humans (H) or animals (A)	Severe disease usually associated with immune suppression	Fatalities reported in humans
<b>Echinococcosis</b>	<i>Echinococcus granulosus</i> , <i>E. multilocularis</i>	contact	dogs, wild canids, cats	H, A		Yes
<b>Ehrlichiosis/ Anaplasmosis</b>	<i>Ehrlichia</i> and <i>Anaplasma</i> species	vector	deer, rodents, horses, dogs	H	✓	Yes
<b>Equine encephalomyelitis (WEE, VEE, EEE)</b>	Togaviridae	vector	birds, horses	H, A (not VEE)		Yes
<b>Erysipeloid</b>	<i>Erysipelothrix rhusiopathiae</i>	contact	Pigs, fish, crustaceans, and mollusks			Yes
<b>Giardiasis</b>	<i>Giardia intestinalis</i> ( <i>G. Lamblia</i> )	contact	thought to be highly species-specific and rarely transmitted from animals to humans	H	✓	No
<b>Hantaviral diseases</b>	Hantaviruses	aerosol	rodents	H		Yes
<b>Herpes B virus</b>	Cercopithecine herpesvirus 1	contact	macaque monkeys			Yes
<b>Influenza</b>	Influenza viruses	contact, aerosol	poultry, swine	H, A		Yes
<b>Larval migrans: cutaneous (hookworm)</b>	<i>Ancylostoma</i> sp.	contact	dogs, cats			Rare
<b>Larval migrans: visceral, ocular, neuro (roundworm)</b>	<i>Toxocara canis</i> , <i>Toxocara cati</i>	contact	dogs, cats			Rare
<b>Leishmaniasis</b>	<i>Leishmania</i> species	vector	dogs, wild canids			Yes
<b>Leptospirosis</b>	<i>Leptospira</i> species	contact, aerosol	rodents, pigs, cattle, sheep, goats, horses, dogs,	A		Yes
<b>Listeriosis</b>	<i>Listeria monocytogenes</i>	contact	cattle, sheep, goats, pigs, birds, dogs, cats	H	✓	Yes
<b>Lyme disease</b>	<i>Borrelia burgdorferi</i>	vector	small rodents, wild mammals	H		No
<b>Lymphocytic choriomeningitis</b>	Arenavirus (LCMV)	contact aerosol	mice, hamsters, guinea pigs		✓	Yes
<b>Monkeypox</b>	Orthopoxvirus	contact, aerosol	non-human primates, rodents	A		Yes
<b>Mycobacteria, non- tuberculous</b>	<i>Mycobacterium avium</i> complex, <i>Mycobacterium marinum</i> ,	aerosol, contact	Poultry, birds, aquarium fish, reptiles		✓	Yes
<b>Pasteurellosis</b>	<i>Pasteurella multocida</i> and other species	contact	dogs, cats, rabbits, rodents		✓	No

## Zoonotic Diseases of Importance in the United States

Disease	Agent	Means of transmission to humans	Most common species associated with transmission to humans	Nationally notifiable for humans (H) or animals (A)	Severe disease usually associated with immune suppression	Fatalities reported in humans
<b>Plague</b>	<i>Yersinia pestis</i>	vector, contact, aerosol	rodents, cats, dogs	H, A		Yes
<b>Psittacosis/Chlamydiosis</b>	<i>Chlamydophila psittaci</i>	aerosol, contact	pet birds, poultry	H	✓	Yes
<b>Q Fever</b>	<i>Coxiella burnetti</i>	contact, aerosol, vector	goats, sheep, cattle, cats, dogs, rodents, lagamorphs	H, A		Yes
<b>Rabies</b>	Lyssavirus	contact	Cats, dogs, cattle and other domestic animals, wild carnivores, raccoons, bats	H, A		Yes
<b>Rat bite fever</b>	<i>Streptobacillus moniliformis</i> , <i>Spirillum minus</i>	contact	rodents		✓	Yes
<b>Rhodococcus equi infection</b>	<i>Rhodococcus equi</i>	aerosol, contact	horses		✓	Yes
<b>Salmonellosis</b>	various <i>Salmonella</i> species	contact	reptiles, poultry, horses, pocket pets, many species of mammals and birds	H	✓	Yes
<b>Sporotrichosis</b>	<i>Sporothrix schenckii</i>	contact	cats, dogs, horses		✓	Rare
<b>Staphylococcosis</b>	<i>Staphylococcus</i> species	contact	dogs, cats, horses	H (MRSA)	✓	Yes (some forms)
<b>Streptococcosis</b>	<i>Streptococcus</i> species	contact, aerosol	swine, other mammals, fish	some forms H		Yes (some forms)
<b>Toxoplasmosis</b>	<i>Toxoplasma gondii</i>	contact	cats		✓	Yes
<b>Trichuriasis (whipworm infection)</b>	<i>Trichuris suis</i> , <i>T. trichiura</i> , <i>T. vulpis</i>	contact	dogs, swine			Rare
<b>Tuberculosis, bovine</b>	<i>Mycobacterium bovis</i>	aerosol, contact	cattle, swine, sheep, goats	H, A		Yes
<b>Tularemia</b>	<i>Francisella tularensis</i>	vector, contact, aerosol	lagamorphs, pocket pets, wild aquatic rodents, sheep, cats, horses, dogs	H, A		Yes
<b>Vesicular stomatitis</b>	Vesicular stomatitis	vector, contact	horses, cattle, swine, sheep, goats	A		No
<b>West Nile Virus</b>	Flaviviridae	vector	wild birds	H		Yes
<b>Yersiniosis</b>	<i>Yersinia enterocolitica</i>	contact	swine, various animal and bird species			No

Note:

-- Nationally reportable diseases were taken from the Centers for Disease Control and Prevention (CDC) nationally notifiable infectious diseases, the World Organization for Animal Health (OIE) notifiable animal diseases, and the USDA high consequence livestock pathogens lists. Cases may also be notifiable at the state level. Check with your State Veterinarian or State Public Health Veterinarian for a current listing of reportable diseases in your area.

References:

Acha PN, Szyfres B. Zoonoses and Communicable Diseases Common to Man and Animals. Volume I. Bacterioses and Mycoses; Volume II. Chlamydioses, Rickettsioses, and Viroses; and Volume III. Parasitoses. Pan American Health Organization: Washington DC 2003.

CDC. Healthy Pets Healthy People website. Available at: [www.cdc.gov/healthypets/browse\\_by\\_diseases.htm](http://www.cdc.gov/healthypets/browse_by_diseases.htm). Accessed Aug 22, 2006.

CDC. Nationally Notifiable Infectious Diseases. United States 2006. Available at: [www.cdc.gov/epo/dphsi/phs/infdis2006.htm](http://www.cdc.gov/epo/dphsi/phs/infdis2006.htm). Accessed on Aug 22, 2006.

Public Health Agency of Canada. Material Safety Data Sheets – Infectious Substances. Available at: [www.phac-aspc.gc.ca/msds-ftss/index.html](http://www.phac-aspc.gc.ca/msds-ftss/index.html). Accessed Aug 22, 2006.

The Merck Veterinary Manual. 9<sup>th</sup> Edition. Kahn CM, Line SL ed. Merck and Company: Whitehouse Station, NJ, 2005.

Trejejo RT, Barr MC, Robinson RA. Important emerging bacterial zoonotic infections affecting the immunocompromised. *Vet Res* 2005;36:493-506.

United States Department of Agriculture. High consequence livestock pathogens list. Available at: [www.ars.usda.gov/SP2UserFiles/Place/54000000/Policies/PM03-003.PDF](http://www.ars.usda.gov/SP2UserFiles/Place/54000000/Policies/PM03-003.PDF). Accessed Aug 22, 2006.

World Organization for Animal Health (OIE). Diseases notifiable to the OIE. Updated 01/23/06. Available at: [www.oie.int/eng/maladies/en\\_classification.htm](http://www.oie.int/eng/maladies/en_classification.htm). Accessed Aug 22, 2006.

## APPENDIX B

### Selected Disinfectants Used in Veterinary Practice

Characteristics of Selected Disinfectants							
Disinfectant Category	Alcohols	Aldehydes	Biguanides	Halogens: Hypochlorites	Halogens: Iodine Compounds	Oxidizing Agents	Phenols
Sample Trade Names	Ethyl alcohol Isopropyl alcohol	Formaldehyde Glutaraldehyde	Chlorhexidine Nolvasan <sup>®</sup> Virosan <sup>®</sup>	Bleach	Betadine <sup>®</sup> Providone <sup>®</sup>	Hydrogen peroxide Peracetic acid Virkon S <sup>®</sup> Oxy-Sept 333 <sup>®</sup>	One-Stroke Environ <sup>®</sup> Pheno-Tek II <sup>®</sup> Tek-Trol <sup>®</sup>
Mechanism of Action	•Precipitates proteins •Denatures lipids	•Denatures proteins •Alkylates nucleic acids	•Alters membrane permeability	•Denatures proteins	•Denatures proteins	•Denature proteins and lipids	•Denatures proteins •Alters cell wall permeability
Advantages	•Fast acting •Leaves no residue	•Broad spectrum	•Broad spectrum	•Broad spectrum •Short contact time •Inexpensive	•Stable in storage •Relatively safe	•Broad spectrum	•Good efficacy with organic material •Non-corrosive •Stable in storage
Disadvantages	•Rapid evaporation •Flammable	•Carcinogenic •Mucous membranes and tissue irritation •Only use in well ventilated areas	•Only functions in limited pH range (5-7) •Toxic to fish (environmental concern)	•Inactivated by sunlight •Requires frequent application •Corrodes metals •Mucous membrane and tissue irritation	•Inactivated by QACs •Requires frequent application •Stains clothes and treated surfaces		•Stable in storage •Non-irritating to skin •Effective at high temperatures and high pH (9-10)
Precautions	Flammable	Carcinogenic		Never mix with acids; toxic chlorine gas will be released			May be toxic to animals, especially cats and pigs
Vegetative Bacteria	Effective	Effective	Effective	Effective	Effective	Effective	Effective
Mycobacteria	Effective	Effective	Variable	Effective	Limited	Effective	Variable
Enveloped Viruses	Effective	Effective	Limited	Effective	Effective	Effective	Effective
Non-enveloped Viruses	Variable	Effective	Limited	Effective	Limited	Effective	Variable
Spores	Not Effective	Effective	Not Effective	Variable	Limited	Variable	Not Effective
Fungi	Effective	Effective	Limited	Effective	Effective	Variable	Variable
Efficacy with Organic Matter	Reduced	Reduced	?	Rapidly reduced	Rapidly reduced	Variable	Inactivated
Efficacy with Hard Water	?	Reduced	?	Effective	?	?	Inactivated
Efficacy with Soap/ Detergents	?	Reduced	Inactivated	Inactivated	Effective	?	Inactivated

DISCLAIMER: The use of trade names does not in any way signify endorsement of a particular product. For additional product names, please consult the most recent Compendium of Veterinary Products.

ADAPTED FROM: Linton AH, Hugo WB, Russel AD. Disinfection in Veterinary and Farm Practice. 1987. Blackwell Scientific Publications: Oxford, England; Quinn PJ, Markey BK. Disinfection and Disease Prevention in Veterinary Medicine. In: Block SS, ed., Disinfection, Sterilization and Preservation. 5th edition. 2001. Lippincott, Williams and Wilkins: Philadelphia.

COSD\_20606

# The Antimicrobial Spectrum of Disinfectants

Chemical Disinfectants  
*Note: Removal of organic material must always precede the use of any disinfectant.*

most susceptible

	Acids (hydrochloric acid, acetic acid, citric acid)	Alcohols (ethyl alcohol, isopropyl alcohol)	Aldehydes (formaldehyde, paraformaldehyde, glutaraldehyde)	Alkalis (sodium or ammonium hydroxide, sodium carbonate)	Biguanides (chlorhexidine*, Nolvasan*, Chlorhex*, Virosan*, Hibistat*)	Halogens hypochlorite iodine	Oxidizing Agents (hydrogen peroxide, peroxyacetic acid, Trilectant*, Virkon-S*, Oxy-Sept 333*)	Phenolic Compounds (Lysol*, Oxy*, Amphyl*, TekTrol*, Pheno-Tek II*)	Quaternary Ammonium Compounds (Roccal*, Zephiran*, Diquat*, Parvosol*, D-256*)
mycoplasmas	+	++	++	++	++	++	++	++	+
gram-positive bacteria	+	++	++	+	++	+	+	++	++
gram-negative bacteria	+	++	++	+	++	+	+	++	+
pseudomonads	+	++	++	+	+	+	+	++	-
rickettsiae	+	+	+	+	+	+	+	+	+
enveloped viruses	+	+	++	+	+	+	+	+	a
chlamydiae	+	+	+	+	+	+	+	+	-
non-enveloped viruses	-	-	+	+	-	+	+	-	-
fungal spores	+	+	+	+	+	+	+	+	+
picornaviruses (i.e. FMD)	+	N	+	+	N	N	+	N	N
parvoviruses	N	N	+	N	N	N	N	N	+
acid-fast bacteria	-	+	+	+	-	+	+	+	-
bacterial spores	+	-	+	+	-	+	b	-	-
coccidia	-	-	-	c	-	-	-	d	-
prions	-	-	-	-	-	-	-	-	-

susceptibility of microorganisms  
to chemical disinfectants

most resistant

LEGEND  
++ highly effective  
+ effective  
+/- limited activity  
- no activity  
N information not available

a-varies with composition  
b-peracetic acid is sporicidal  
c-ammonium hydroxide  
d-some have activity against coccidia



the Center for  
Food Security  
& Public Health  
IOWA STATE UNIVERSITY®  
www.cfsph.iastate.edu  
AS00\_20606

DISCLAIMER: The use of trade names does not in any way signify endorsement of a particular product. For additional product names, please consult the most recent Compendium of Veterinary Products. ADAPTED FROM: Linton AH, Hugo WB, Russel AD. Disinfection in Veterinary and Farm Practice. 1987. Blackwell Scientific Publications; Oxford, England; Quinn PJ, Markey BK. Disinfection and Disease Prevention in Veterinary Medicine, In: Block SS, ed., Disinfection, Sterilization and Preservation. 5th edition. 2001. Lippincott, Williams and Wilkins: Philadelphia.



## **APPENDIX C**

### **OSHA Respiratory Protection Standard**

Sec. 1910.134 of Appendix D (Mandatory) Information for Employees Using Respirators When Not Required Under the Standard

Respirators are an effective method of protection against designated hazards when properly selected and worn. Respirator use is encouraged, even when exposures are below the exposure limit, to provide an additional level of comfort and protection for workers. However, if a respirator is used improperly or not kept clean, the respirator itself can become a hazard to the worker. Sometimes, workers may wear respirators to avoid exposures to hazards, even if the amount of hazardous substance does not exceed the limits set by OSHA standards. If your employer provides respirators for your voluntary use, or if you provide your own respirator, you need to take certain precautions to be sure that the respirator itself does not present a hazard.

You should do the following:

1. Read and heed all instructions provided by the manufacturer on use, maintenance, cleaning and care, and warnings regarding the respirator's limitations.
2. Choose respirators certified for use to protect against the contaminant of concern. NIOSH, the National Institute for Occupational Safety and Health of the U.S. Department of Health and Human Services, certifies respirators. A label or statement of certification should appear on the respirator or respirator packaging. It will tell you what the respirator is designed for and how much it will protect you.
3. Do not wear your respirator into atmospheres containing contaminants for which your respirator is not designed to protect against. For example, a respirator designed to filter dust particles will not protect you against gases, vapors, or very small solid particles of fumes or smoke.
4. Keep track of your respirator so that you do not mistakenly use someone else's respirator.

## APPENDIX D

### Model Infection Control Plan for Veterinary Practices, 2006

#### National Association of Public Health Veterinarians (NASPHV) Veterinary Infection Control Committee (VICC)

*This model plan should be adapted to your practice in keeping with local, state and federal regulations. A modifiable electronic version is available on the website of the National Association of State Public Health Veterinarians at [www.nasphv.org](http://www.nasphv.org). Please refer the corresponding sections in the full Compendium of Veterinary Standard Precautions for complete information and guidance (also available at [www.nasphv.org](http://www.nasphv.org)).*

Clinic: \_\_\_\_\_

Date of Plan Adoption: \_\_\_\_\_

Date of Next Review: \_\_\_\_\_

Infection Control Officer: \_\_\_\_\_

This plan will be followed as part of our clinic's routine practices. The plan will be reviewed at least annually and as part of new employee training.

#### PERSONAL PROTECTIVE ACTIONS AND EQUIPMENT

**Hand Hygiene:** Wash hands before and after each patient encounter and after contact with blood, body fluids, secretions, excretions or articles contaminated by these fluids. Wash hands before eating, drinking or smoking; after using the toilet; after cleaning animal cages or animal care areas; and whenever hands are visibly soiled. Alcohol-based gels may be used if hands are not visibly soiled, but handwashing with soap and running water is preferred. Keep fingernails short. Keep handwashing supplies stocked at all times. Staff responsible \_\_\_\_\_.

##### **Correct handwashing procedure:**

- Wet hands with running water
- Place soap in palms
- Rub hands together to make a lather
- Scrub hands vigorously for 20 seconds
- Dry hands with a disposable towel
- Turn off faucet handle using the disposable towel

**Use of Gloves and Sleeves:** Wear gloves or sleeves when touching blood, body fluids, secretions, excretions, mucous membranes, and non-intact skin. Wear gloves for dentistry, resuscitations, necropsies, and obstetrical procedures; when cleaning cages and contaminated environmental surfaces and equipment; when handling dirty laundry; when

handling diagnostic specimens (e.g., urine, feces, aspirates, swabs); and when handling an animal with a suspected infectious disease. Change gloves between examination of individual animals or animal groups (e.g., a litter of puppies) and between dirty and clean procedures on the same patient. Gloves should be removed promptly and disposed of after use. Disposable gloves should not be washed and reused. Hands should be washed immediately after glove removal.

Note: Gloves are not necessary when examining or handling normal, healthy animals.

**Facial Protection:** Wear facial protection whenever exposure to splashes or sprays is likely to occur. Facial protection should include a mask worn with either goggles or a face shield. Wear facial protection for the following procedures: dentistry, resuscitation, nebulization, suctioning, bronchoscopy, wound irrigation, obstetrical procedures, and necropsies. Use a surgical mask when cleaning with high-pressure sprayers.

**Respiratory Protection:** Wear a disposable N-95 respirator or other particulate respirator when investigating abortions in small ruminants or significant poultry mortality, when handling ill psittacine birds, and in any other circumstance where there is concern about aerosol transmission.

**Protective Outerwear:** Wear a protective outer garment such as a lab coat, smock, non-sterile gown, or coveralls when attending animals and when conducting cleaning chores. These should be changed whenever soiled, after handling an animal with a known or suspected infectious disease, after working in the isolation room, and after performing a necropsy or other high-risk procedure. Shoes or boots should have thick soles and closed toes, and be water resistant and easily cleanable. Disposable shoe covers should be worn when heavy quantities of infectious materials are present or expected. Impermeable outwear should be worn during obstetrical procedures and necropsies and whenever substantial splashes or large quantities of body fluids may be encountered. Keep clean outer garments available at all times. Staff responsible \_\_\_\_\_.

**Bite and Other Animal-Related Injury Prevention:** Take precautions to prevent bites and other injuries. Use physical restraints, muzzles, bite-resistant gloves, and sedation or anesthesia as needed. Plan an escape route when handling large animals. Do not rely on owners or untrained staff for animal restraint. Notify \_\_\_\_\_ if there concern for personal safety. When bites or scratches occur, wash the site with soap and water immediately. Report all bites and other injuries to \_\_\_\_\_ (Infection Control Officer) who will also maintain the incident report log. Consult a physician whenever the skin is broken. An evaluation of the need for medical attention, tetanus immunization, antibiotics, and rabies post-exposure prophylaxis will be made by a physician. Bite incidents will be reported to \_\_\_\_\_ (public health agency) as required by law. Telephone number: \_\_\_\_\_.

## PROTECTIVE ACTIONS DURING VETERINARY PROCEDURES

**Intake:** Avoid bringing aggressive or potentially infectious animals in through the reception area. If they must come through the main entrance, carry the animal or place it on a gurney so that it can be taken directly into an exam room.

**Examination of Animals:** Wear appropriate protective outward and wash hands before and after examination of individual animals or animal groups (e.g., a litter of puppies). Potentially infectious animals will be examined in a dedicated exam room and remain there until diagnostic procedures and treatments have been performed.

**Injections, Venipuncture, and Aspirations:** Wear gloves while performing venipuncture on animals suspected of having an infectious disease and when performing soft tissue aspirations. Currently, there is no data indicating that venipuncture on healthy animals carries a significant risk of infection.

**Needlestick Injury Prevention:** Do not recap needles except in rare instances when required as part of a medical procedure or protocol. Dispose of all sharps in designated puncture-proof sharps containers. Dispose of the used syringe with attached needle in the sharps container when injecting live vaccines or aspirating body fluids. For most other veterinary procedures, use the needle removal device on the sharps container and dispose of the syringe in the regular trash. Sharps containers are located in every area of the clinic where sharps are used. Do not transfer sharps from one container to another.

**Dental Procedures:** Wear protective outerwear, gloves, mask, and a face shield or goggles when performing dental procedures or working nearby (such as when monitoring anesthesia).

**Resuscitation:** Wear gloves, mask, and a face shield or goggles.

**Obstetrics:** Wear gloves and/or shoulder-length sleeves, mask or respirator, face shield or goggles, and impermeable outerwear.

**Necropsy:** Wear cut-resistant gloves, mask, face shield or goggles, and impermeable outerwear. Only necessary personnel are allowed in the vicinity of the procedure. Wear a respirator when using a band saw or other power equipment. If an animal is suspected of having a notifiable infectious or a foreign animal disease, consult with the State Veterinarian before proceeding with a necropsy. Contact information for State Veterinarian's office\_\_\_\_\_.

**Diagnostic Specimen Handling:** Wear protective outerwear and gloves. Discard gloves and wash hands before touching clean items (e.g., microscope, telephone). Eating and drinking are not allowed in the laboratory.

## ENVIRONMENTAL INFECTION CONTROL

**Isolation of Infectious Animals:** Animals with a contagious or zoonotic disease will be housed in isolation as soon as possible. Clearly mark the room or cage to indicate the patient's status and describe additional precautions. Only equipment needed for the care and treatment of the patient should be kept in the isolation room, and there should also be dedicated cleaning supplies. Disassemble and thoroughly clean and disinfect any equipment that must be taken out of the room. Discard gloves after use. Leave other personal protective equipment (e.g., gown, mask) in the isolation room for reuse. Clean and disinfect protective equipment between patients and whenever contaminated by body fluids. Bag potentially contaminated materials before removal from the isolation room. Use disinfectant footbath before entering and leaving the room. Access to the isolation room is limited. Keep a sign-in log of all people (including owners or other non-employees) having contact with a patient in isolation. Monitor air pressure daily while the room is in use.

Staff responsible \_\_\_\_\_

**Cleaning and Disinfection of Equipment and Environmental Surfaces:** Clean surfaces and equipment first to remove organic matter, and then use an EPA-registered hospital disinfectant, applied according to manufacturer's instructions. Minimize dust and aerosols when cleaning. Clean and disinfect animal cages, toys, and food and water bowls between animals and whenever visibly soiled. Clean litter boxes once a day. Wear gloves when cleaning, and wash hands afterwards. There is a written checklist for each area of the facility (e.g., waiting room, exam rooms, treatment area, kennels) specifying the frequency of cleaning, disinfection procedures, products to be used, and staff responsible.

**Handling Laundry:** Wear gloves when handling soiled laundry. Wash animal bedding and other laundry with standard laundry detergent and machine dry. Use separate storage and transport bins for clean and dirty laundry.

**Decontamination and Spill Response:** Immediately spray a spill or splash of blood or body fluids with disinfectant and contain it with absorbent material (e.g., paper towels, sawdust, cat litter). Don gloves, mask, and protective clothing (including shoe covers if the spill is on the floor and may be stepped in) before beginning the clean-up. Pick up the material then seal in leak-proof plastic bags. Clean the area, and disinfect according to manufacturer's instructions. Keep clients, patients and employees away from the spill area until disinfection is completed.

**Veterinary Medical Waste:** *Insert here your local and state ordinances regulating disposal of animal waste, pathology waste, animal carcasses, bedding, sharps and biologics.*

**Rodent and Vector Control:** Keep the facility free of wild rodents and mosquitoes and other arthropods by sealing entry portals, eliminating clutter and sources of standing water, keeping animal food in closed metal or thick plastic containers, and disposing of food waste properly.

**Other Environmental Controls:** There are designated areas for eating, drinking, smoking, applying make-up and similar activities. These activities should never be done in animal care areas or in the laboratory area. Human food or drink should not be kept in the same refrigerator as animal food, biologics, or laboratory specimens. Dishes for human use should be cleaned and stored away from animal care and animal food preparation areas.

## **EMPLOYEE HEALTH**

The following personnel are responsible for developing and maintaining the practice's infection control policies, keeping records, and managing workplace exposure and injury incidents.

Staff Responsible:

---

---

---

### **Employee Immunization Policies and Record Keeping:**

**Record Keeping:** Current emergency contact information will be maintained for each employee. Records will be maintained on immunizations, rabies titers, and exposure and injury incidents. Report and record changes in health status (e.g. pregnancy) that may affect work duties.

**Rabies Preexposure Vaccination:** All staff with animal contact must be vaccinated against rabies, followed by periodic titer checks and rabies boosters, in accordance with the recommendations of the Advisory Committee on Immunization Practices (CDC, 1999).

**Tetanus Vaccination:** Tetanus immunizations must be up to date. Report and record puncture wounds and other possible exposures to tetanus. Consult a health care provider regarding the need for a tetanus booster.

**Seasonal Influenza Vaccination:** Unless contraindicated, veterinary personnel are encouraged to receive the current seasonal influenza vaccine. *Check with CDC for updated recommendations at [www.cdc.gov](http://www.cdc.gov).*

**Staff Training and Education:** Infection control training and education will be documented in the employee health record.

**Documenting and Reporting Exposure Incidents:** Report incidents that result in injury or potential exposure to an infectious agent to \_\_\_\_\_.

The following information will be collected for each exposure incident: date, time, location, person(s) injured or exposed, other persons present, description of the incident, the status of any animals involved (e.g., vaccination history, clinical condition, diagnostic

information), and plans for follow-up. If consultation with a health care provider is necessary, be sure to inform them of the exposure to the animal(s).

**Pregnant and Immunocompromised Personnel:** Pregnant and immunocompromised employees are at increased risk from zoonotic diseases. Inform \_\_\_\_\_ if you are concerned about your work responsibilities, so that accommodations may be made. Consultation between the supervising veterinarian and a health care provider may be needed.

**The following information is attached to the Infection Control Plan:**

- List of reportable/notifiable veterinary diseases and where to report
- State and local public health contacts for consultation on zoonotic diseases
- Public Health Laboratory services and contact information
- Emergency services telephone numbers – fire, police, sheriff, animal control, poison control, etc
- List of EPA-registered disinfectants
- OSHA regulations
- State Department of Agriculture or Board of Animal Health contact information and regulations
- Local animal waste disposal and biohazard regulations
- Local rabies regulations
- Local animal control and exotic animal regulations
- Useful resources